INDEX TO AUTHORS

BECK, S., see *Peacock, P. R.*

BERENBLUM, I., and SCHOENTAL, R., carcinogenic constituents of shale oil, 232

BERGER, F. M., the sensitizing and indicator action of Victoria blue and Janus green on the flocculation reaction for syphilis, 252

BERNKOPF, H., see *Kligler, I. J.*

CARR, J. G., prolonged antibody production following recovery of fowls from Rous No. 1 sarcoma, 138

——— some investigations upon the nature of the resistance of an inbred line of fowls to the development of the Rous No. 1 sarcoma, 127

——— the relation between age, structure, and agent content of Rous No. 1 sarcoma, 133

CHAIN, E., FLOREY, H. W., JENNINGS, M. A., and WILLIAMS, T. I., helvolic acid, an antibiotic produced by *Aspergillus fumigatus, mut. helvola* Yuill, 108

CROWFOOT, D. M., and LOW, B. W., a note on the crystallography of helvolic acid and the methyl ester of helvolic acid, 120

DANIELLI, J. F., see *Fell, H. B.*

DMOCHOWSKI, L., and GYE, W. E., influence of cross-suckling on the incidence of mammary cancer in high and low-cancer strains of mice, 223

DONIACH, I., MOTTRAM, J. C., and WEIGERT, F., the fluorescence of 3 : 4-benzpyrene *in vivo*.
I. The distribution of fluorescence at various sites, especially the skin of mice, 1
II. The inter-relationship of the derivatives formed in various sites, 9

ELLIOTT, S. D., type relationships amongst group A streptococci, 159

EVANS, D. G., the protective properties of the alpha antitoxin and theta antihaemolysin occurring in *Cl. welchii* type A antiserum, 81

FELL, H. B., and DANIELLI, J. F., the enzymes of healing wounds. I. The distribution of alkaline phosphomonoesterase in experimental wounds and burns in the rat, 196

FINDLAY, G. M., see *Maegraith, B. G.*

FLOREY, H. W., see *Chain, E.*

FULTON, F., staphylococcal enterotoxin, with special reference to the kitten test, 65

GRIESBACH, W. E., and PURVES, H. D., studies on experimental goitre. V. Pituitary function in relation to goitrogenesis and thyroidectomy, 174

——— ——— the assay of thyrotropic activity by the cell height response in guinea-pigs, 185

GYE, W. E., see *Dmochowski, L.*

HARRIS, G. C. M., see *Wilkins, W. H.*

JACKSON, E. B., see *Turner, J. C.*

JAPA, J., a study of the morphology and development of the megakaryocytes, 73

JENNINGS, M. A., see *Chain, E.*

KASSANIS, B., neutralization of some plant viruses by rabbit sera, 152

KEELE, C. A., see *Scarff, R. W.*

KLIGLER, I. J., and BERNKOPF, H., the path of dissemination of rabies virus in the body of normal and immunized mice, 15

LOW, B. W., see *Crowfoot, D. M.*

MCILWAIN, H., nutritional studies of bacterial variation.
I. Resistance to pantoyltaurine in naturally-occurring and experimentally prepared strains, 203
II. The derivation of drug-resistant strains in the absence of any inhibitor, 212

MCINTOSH, J., and SELBIE, F. R., the production of drug resistant cultures of bacteria *in vitro* and a study of their inter-relationships, 246

MACOUN, S. J. R., see *Prunty, F. T. G.*

MAEGRAITH, B. G., MARTIN, N. H., and FINDLAY, G. M., the mechanism of red blood cell destruction, 58

MARTIN, N. H., see *Maegraith, B. G.*

MERLING, K. B. E., ultramicroscopical observations on the morphology and development of vaccinia virus *in vitro*, 240

MILES, A. A., and MILES, E. M., the fixation of foreign material in inflamed tissue, with especial reference to the action of *Cl. welchii* toxin and antitoxin, 95

MILES, E. M., see *Miles, A. A.*

MORGAN, W. T. J., an artificial antigen with blood-group A specificity, 41

MOTTRAM, J. C., see *Doniach, I.*

OSBORN, E. M., on the occurrence of antibacterial substances in green plants, 227

PAGE, J. E., and ROBINSON, F. A., an examination of the relationship between the bacteriostatic activity and the normal reduction potentials of substituted quinones, 89
PEACOCK, P. R., and BECK, S., multiple mesenteric sarcomata in rats following ingestion of heated lard, 143
PRUNTY, F. T. G., and MACOUN, S. J. R., the chloride excretion in steatorrhoea: comparison with the conditions in Addison's disease, 22
PURVES, H. D., studies on experimental goitre. IV. The effect of di-iodotyrosine and thyroxine on the goitrogenic action of brassica seeds, 171
——— see *Griesbach, W. E.*

RENNIE, J. B., a modified laevulose (fructose) tolerance test compared with the hippuric acid synthesis test of hepatic efficiency, 26
ROBINSON, F. A., see *Page, J. E.*
ROBSON, J. M., and SCOTT, G. I., the production and treatment of experimental pneumococcal hypopyon ulcers in the rabbit, 50

SCARFF, R. W., and KEELE, C. A., the effects of temporary occlusion of the renal circulation in the rabbit, 147
SCHOENTAL, R., see *Berenblum, I.*
SCOTT, G. I., see *Robson, J. M.*
SELBIE, F. R., viability of *Treponema pallidum* in stored plasma, 150
——— see *McIntosh, J.*
SIMPSON, R. E. H., criteria of life and death in human pus cells, 218
STUART-HARRIS, C. H., observations on the agglutination of fowl red cells by influenza viruses, 33

TENENBAUM, E., a pure strain of Rous sarcoma cells, 56
TURNER, J. C., and JACKSON, E. B., serological specificity of an autoantibody in atypical pneumonia, 121

WEIGERT, F., see *Doniach, I.*
WHITEHEAD, V. I. E., basal metabolism of the albino rat fed on goitrogenic diet, 192
WILKINS, W. H., and HARRIS, G. C. M., investigation into the production of bacteriostatic substances by fungi. Preliminary examination of a second 100 fungal species, 141
WILLIAMS, T. I., see *Chain, E.*